

## Publication list Anne Thorup, Feb 2017

Submitted:

\*Thorup A, Laursen, TM, Munk-Olsen, T, Ranning A, Mortensen PB, Plessen K, Nordentoft M ' Incidence of child and adolescent mental illness in children aged 0-17 with familial high risk for severe mental illness – a Danish Register Study'

In press:

Burton BK, Thorup A.A.E., Jepsen JR, Poulsen G, Ellersgaard D, Spang KS, Christiani CJ, Hemager N, Gantriis D, Greve A, Mors O, Nordentoft M, Plessen KJ (2016) Impairments of Motor Function among Children with a Familial Risk of Schizophrenia or Bipolar Disorder at 7 Years Old – an observational cohort study. *Lancet Psychiatry* (in press)

### 2016

Ranning A, Munk LT, Thorup A, et al. Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. *J Am Acad Child Adolesc Psychiatry* 2016;55:953-61

Kuhl JO, Laursen TM, Thorup A, Nordentoft M. The incidence of schizophrenia and schizophrenia spectrum disorders in Denmark in the period 2000-2012. A register-based study. *Schizophr.Res.* 2016;

Burton BK, Hjorthoj C, Jepsen JR, Thorup A, Nordentoft M, Plessen KJ. Research Review: Do motor deficits during development represent an endophenotype for schizophrenia? A meta-analysis. *J. Child Psychol. Psychiatry* 2016; 57:446-456

### 2015

Melau M, Harder S, Jeppesen P, Hjorthoj C, Jepsen JR, Thorup A, Nordentoft M. The association between working alliance and clinical and functional outcome in a cohort of 400 patients with first-episode psychosis: a cross-sectional study. *J.Clin.Psychiatry* 2015; 76:e83-e90

\*Thorup AA, Jepsen JR, Ellersgaard DV, Burton BK, Christiani CJ, Hemager N, Skjaerbaek M, Ranning A, Spang KS, Gantriis DL, Greve AN, Zahle KK, Mors O, Plessen KJ, Nordentoft M. The Danish High Risk and Resilience Study - VIA 7 - a cohort study of 520 7-year-old children born of parents diagnosed with either schizophrenia, bipolar disorder or neither of these two mental disorders. *BMC.Psychiatry* 2015; 15:233

Nordentoft M, Melau M, Iversen T, Petersen L, Jeppesen P, Thorup A, Bertelsen M, Hjorthoj CR, Hastrup LH, Jorgensen P. From research to practice: how OPUS treatment was accepted and implemented throughout Denmark. *Early Interv.Psychiatry* 2015; 9:156-162

Austin SF, Mors O, Budtz-Jorgensen E, Secher RG, Hjorthoj CR, Bertelsen M, Jeppesen P, Petersen L, Thorup A, Nordentoft M. Long-term trajectories of positive and negative symptoms in first episode psychosis: A 10year follow-up study in the OPUS cohort. *Schizophr.Res.* 2015;

Burton BK, Hjorthoj C, Jepsen JR, Thorup A, Nordentoft M, Plessen KJ. Research Review: Do motor deficits during development represent an endophenotype for schizophrenia? A meta-analysis. *J.Child Psychol.Psychiatry* 2015;

Ranning A, Munk LT, Thorup A, Hjorthoj C, Nordentoft M. Serious mental illness and disrupted caregiving for children: a nationwide, register-based cohort study. *J.Clin.Psychiatry* 2015; 76:e1006-e1014

Secher RG, Hjorthoj CR, Austin SF, Thorup A, Jeppesen P, Mors O, Nordentoft M. Ten-year follow-up of the OPUS specialized early intervention trial for patients with a first episode of psychosis. *Schizophr.Bull.* 2015; 41:617-626

## **2014**

Clausen L, Hjorthoj CR, Thorup A, Jeppesen P, Petersen L, Bertelsen M, Nordentoft M. Change in cannabis use, clinical symptoms and social functioning among patients with first-episode psychosis: a 5-year follow-up study of patients in the OPUS trial. *Psychol.Med.* 2014; 44:117-126

\*Thorup A, Albert N, Bertelsen M, Petersen L, Jeppesen P, Le QP, Krarup G, Jorgensen P, Nordentoft M. Gender differences in first-episode psychosis at 5-year follow-up--two different courses of disease? Results from the OPUS study at 5-year follow-up. *Eur.Psychiatry* 2014; 29:44-51

Nordentoft M, Rasmussen JO, Melau M, Hjorthoj CR, Thorup AA. How successful are first episode programs? A review of the evidence for specialized assertive early intervention. *Curr.Opin.Psychiatry* 2014; 27:167-172

Ellersgaard D, Mors O, Thorup A, Jorgensen P, Jeppesen P, Nordentoft M. Prospective study of the course of delusional themes in first-episode non-affective psychosis. *Early Interv.Psychiatry* 2014; 8:340-347

## **2013**

Hastrup LH, Kronborg C, Bertelsen M, Jeppesen P, Jorgensen P, Petersen L, Thorup A, Simonsen E, Nordentoft M. Cost-effectiveness of early intervention in first-episode psychosis: economic evaluation of a randomised controlled trial (the OPUS study). *Br.J.Psychiatry* 2013; 202:35-41

Nordentoft M, Melau M, Iversen T, Petersen L, Jeppesen P, Thorup A, Bertelsen M, Hjorthoj CR, Hastrup LH, Jorgensen P. From research to practice: how OPUS treatment was accepted and implemented throughout Denmark. *Early Interv.Psychiatry* 2013;

Austin SF, Mors O, Secher RG, Hjorthoj CR, Albert N, Bertelsen M, Jensen H, Jeppesen P, Petersen L, Randers L, Thorup A, Nordentoft M. Predictors of recovery in first episode psychosis: the OPUS cohort at 10 year follow-up. *Schizophr.Res.* 2013; 150:163-168

## 2011

Melau M, Jeppesen P, Thorup A, Bertelsen M, Petersen L, Gluud C, Krarup G, Nordentoft M. The effect of five years versus two years of specialised assertive intervention for first episode psychosis - OPUS II: study protocol for a randomized controlled trial. *Trials* 2011; 12:72

Albert N, Bertelsen M, Thorup A, Petersen L, Jeppesen P, Le Quack P, Krarup G, Jorgensen P, Nordentoft M. Predictors of recovery from psychosis Analyses of clinical and social factors associated with recovery among patients with first-episode psychosis after 5years. *Schizophr.Res.* 2011; 125:257-266

## 2010

Nordentoft M, Ohlenschlaeger J, Thorup A, Petersen L, Jeppesen P, Bertelsen M. Deinstitutionalization revisited: a 5-year follow-up of a randomized clinical trial of hospital-based rehabilitation versus specialized assertive intervention (OPUS) versus standard treatment for patients with first-episode schizophrenia spectrum disorders. *Psychol.Med.* 2010; 40:1619-1626

\*Thorup A, Petersen L, Jeppesen P, Nordentoft M. The quality of life among first-episode psychotic patients in the OPUS trial. *Schizophr.Res.* 2010; 116:27-34

## 2009

Bertelsen M, Jeppesen P, Petersen L, Thorup A, Ohlenschlaeger J, Le Quach P, Ostergaard CT, Krarup G, Jorgensen P, Nordentoft M. Course of illness in a sample of 265 patients with first-episode psychosis--five-year follow-up of the Danish OPUS trial. *Schizophr.Res.* 2009; 107:173-178

Bertelsen M, Jeppesen P, Petersen L, Thorup A, Ohlenschlaeger J, Le Quach P, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. [First episode of psychosis intensive early intervention programme versus standard treatment--secondary publication]. *Ugeskr.Laeger* 2009; 171:2992-2995

Quach PL, Mors O, Christensen TO, Krarup G, Jorgensen P, Bertelsen M, Jeppesen P, Petersen L, Thorup A, Nordentoft M. Predictors of poor adherence to medication among patients with first-episode schizophrenia-spectrum disorder. *Early Interv.Psychiatry* 2009; 3:66-74

Nordentoft M, Jeppesen P, Petersen L, Bertelsen M, Thorup A. The rationale for early intervention in schizophrenia and related disorders. *Early Interv.Psychiatry* 2009; 3 Suppl 1:S3-S7

## 2008

Jeppesen P, Petersen L, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. The association between pre-morbid adjustment, duration of untreated psychosis and outcome in first-episode psychosis. *Psychol.Med.* 2008; 38:1157-1166

Nordentoft M, Larsen T, Bertelsen M, Thorup A. The challenge of suicidal behaviour in early psychosis. *Early Interv.Psychiatry* 2008; 2:1-2

Nordentoft M, Thorup A, Petersen L, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Jeppesen P. Does a detection team shorten duration of untreated psychosis? *Early Interv.Psychiatry* 2008; 2:22-26

Ohlenschlaeger J, Nordentoft M, Thorup A, Jeppesen P, Petersen L, Christensen TO, Krarup G, Jorgensen P. Effect of integrated treatment on the use of coercive measures in first-episode schizophrenia-spectrum disorder. A randomized clinical trial. *Int.J.Law Psychiatry* 2008; 31:72-76

Bertelsen M, Jeppesen P, Petersen L, Thorup A, Ohlenschlaeger J, Le Quach P, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. Five-year follow-up of a randomized multicenter trial of intensive early intervention vs standard treatment for patients with a first episode of psychotic illness: the OPUS trial. *Arch.Gen.Psychiatry* 2008; 65:762-771

Nordentoft M, Bertelsen M, Petersen L, Jeppesen P, Thorup A. [Phases of psychosis]. *Ugeskr.Laeger* 2008; 170:3746-3748

Petersen L, Thorup A, Ohlenschlaeger J, Christensen TO, Jeppesen P, Krarup G, Jorgensen P, Mortensen EL, Nordentoft M. Predictors of remission and recovery in a first-episode schizophrenia spectrum disorder sample: 2-year follow-up of the OPUS trial. *Can.J.Psychiatry* 2008; 53:660-670

## **2007**

\*Thorup A, Petersen L, Jeppesen P, Nordentoft M. Frequency and predictive values of first rank symptoms at baseline among 362 young adult patients with first-episode schizophrenia Results from the Danish OPUS study. *Schizophr.Res.* 2007;

\*Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Gender differences in young adults with first-episode schizophrenia spectrum disorders at baseline in the Danish OPUS study. *J.Nerv.Ment.Dis.* 2007; 195:396-405

Ohlenschlaeger J, Thorup A, Petersen L, Jeppesen P, Koster A, Munkner R, Nordentoft M. Intensive treatment models and coercion. *Nord.J.Psychiatry* 2007; 61:369-378

Petersen L, Jeppesen P, Thorup A, Ohlenschlaeger J, Krarup G, Ostergard T, Jorgensen P, Nordentoft M. Substance abuse and first-episode schizophrenia-spectrum disorders. The Danish OPUS trial. *Early Interv.Psychiatry* 2007; 1:88-96

Bertelsen M, Jeppesen P, Petersen L, Thorup A, Ohlenschlaeger J, Le Quach P, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. Suicidal behaviour and mortality in first-episode psychosis: the OPUS trial. *Br.J.Psychiatry Suppl* 2007; 51:s140-s146

\*Thorup A, Waltoft BL, Pedersen CB, Mortensen PB, Nordentoft M. Young males have a higher risk of developing schizophrenia: a Danish register study. *Psychol.Med.* 2007; 37:479-484

## 2006

Nordentoft M, Petersen L, Jeppesen P, Thorup AA, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P. [OPUS: a randomised multicenter trial of integrated versus standard treatment for patients with a first-episode psychosis--secondary publication]. *Ugeskr.Laeger* 2006; 168:381-384

\*Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Social network among young adults with first-episode schizophrenia spectrum disorders: results from the Danish OPUS trial. *Soc.Psychiatry Psychiatr.Epidemiol.* 2006; 41:761-770

Nordentoft M, Thorup A, Petersen L, Ohlenschlaeger J, Melau M, Christensen TO, Krarup G, Jorgensen P, Jeppesen P. Transition rates from schizotypal disorder to psychotic disorder for first-contact patients included in the OPUS trial. A randomized clinical trial of integrated treatment and standard treatment. *Schizophr.Res.* 2006; 83:29-40

## 2005

Petersen L, Nordentoft M, Jeppesen P, Ohlenschlaeger J, Thorup A, Christensen TO, Krarup G, Dahlstrom J, Haastруп B, Jorgensen P. Improving 1-year outcome in first-episode psychosis: OPUS trial. *Br.J.Psychiatry Suppl* 2005; 48:s98-103

\*Thorup A, Petersen L, Jeppesen P, Ohlenschlaeger J, Christensen T, Krarup G, Jorgensen P, Nordentoft M. Integrated treatment ameliorates negative symptoms in first episode psychosis-results from the Danish OPUS trial. *Schizophr.Res.* 2005; 79:95-105

Jeppesen P, Petersen L, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Hemmingsen R, Jorgensen P, Nordentoft M. Integrated treatment of first-episode psychosis: effect of treatment on family burden: OPUS trial. *Br.J.Psychiatry Suppl* 2005; 48:s85-s90

Petersen L, Jeppesen P, Thorup A, Abel MB, Ohlenschlaeger J, Christensen TO, Krarup G, Jorgensen P, Nordentoft M. A randomised multicentre trial of integrated versus standard treatment for patients with a first episode of psychotic illness. *BMJ* 2005; 331:602

1. Ranning A, Munk LT, Thorup A, et al. Children of Parents With Serious Mental Illness: With Whom Do They Grow Up? A Prospective, Population-Based Study. *J Am Acad Child Adolesc Psychiatry* 2016;55:953-61